

KC600 Anti Human Prolyl 4-Hydroxylated FGA Monoclonal Antibody
(Clone No. 11A5)

This product is generated from GANP®




Code No.	KC600
Category	Cancer
Target	α -Fibrinogen with hydroxylation of its proline 565 residue
Type	Monoclonal Antibody
Concentration	0.25mg/ml
Contents (Volume)	50 μ g (200 μ L/vial)
Gene ID	64435
Primary Source	FGA
Synonyms	Fib2; MGC119422; MGC119423; MGC119425; FGA
Immunogen	Partial peptide of Human FGA (ESSSHHP(0)GIAEFPSR [P(0): hydroxyproline])
Raised in	GANP® mouse
Myeloma	P3U1
Clone number	11A5
Purification	
Source	Serum-free medium
Isotype	IgG1, κ
Cross Reactivity	Not Tested
Label	Unlabeled
Buffer	PBS [containing 2% Block Ace as a stabilizer, 0.1% Proclin as a bacteriostat]
Storage	Store below -20°C. Once thawed, store at 4°C. Repeated freeze-thaw cycles should be avoided.
Application	ELISA, WB

Recommended Antibody Dilutions

ELISA	WB	HC	IC
1.0	2.0	Not Tested	Not Tested
P	FCM	F	Neutralization
Not Tested	Not Tested	Not Tested	Not Tested

(μ g/ml)



[WB] Immunoblot analysis of plasma samples from pancreatic cancer patients and healthy controls with 11A5 (upper panel) and anti-fibrinogen (lower panel) antibodies.

UniPlot Summary	<p>//Function: Fibrinogen has a double function: yielding monomers that polymerize into fibrin and acting as a cofactor in platelet aggregation.</p> <p>//Subcellular location: Secreted.</p> <p>//Tissue specificity: Plasma.</p> <p>//Sequence similarities: Contains 1 fibrinogen C-terminal domain.</p>
Reference	<p>1) I. Ono M, et al. "Prolyl 4-Hydroxylation of α-Fibrinogen" J. Biol. Chem. 2009 Oct;284(42):29041-9.</p>